

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
4 July 2002 (04.07.2002)

PCT

(10) International Publication Number
WO 02/051985 A2

(51) International Patent Classification⁷: C12N 5/00

(74) Agent: ATKINSON, Peter, Birch; Marks & Clerk, Sussex House, 83-85 Mosley Street, Manchester M2 3LG (GB).

(21) International Application Number: PCT/GB01/05606

(22) International Filing Date:
19 December 2001 (19.12.2001)

(25) Filing Language: English

(26) Publication Language: English

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(30) Priority Data:
0031462.5 22 December 2000 (22.12.2000) GB
0031651.3 23 December 2000 (23.12.2000) GB
60/267,032 7 February 2001 (07.02.2001) US

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): KEELE UNIVERSITY [GB/GB]; Keele, Staffordshire ST5 5BG (GB).

Published:

— without international search report and to be republished upon receipt of that report

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): EL-HAJ, Alicia, Jennifer, Hafeeza [GB]; Beardhough Cottage, Whitle Road, New Mills, High Peak SK22 4BQ (GB). DOBSON, Jon, Paul [GB]; Claremount Brook Lane, Endon, stoke-on-Trent ST9 9EZ (GB).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 02/051985 A2

(54) Title: CULTURING TISSUE USING MAGNETICALLY GENERATED MECHANICAL STRESSES

(57) Abstract: A method of culturing tissue comprises growing tissue forming cells whilst subjecting the tissue forming cells to mechanical stresses which are generated magnetically.

CULTURING TISSUE USING MAGNETICALLY GENERATED MECHANICAL STRESSES

The present invention relates to a method of culturing cells and relates more particularly (but not exclusively) to the culturing of cells to form replacement human or animal tissue. The invention relates even more particularly, but again not exclusively, to the culturing of mechano-responsive tissue.

The *in vitro* cultivation of replacement tissue for humans and animals is an important development allowing the tissue to be grown from cells taken from the patient so that the replacement tissue does not cause rejection problems. Examples of replacement tissues that may be produced for such replacement therapies include connective tissue, bone, cartilage, tendon and pancreas.

The replacement tissue must not only be comprised of the same type of cells as the tissue it is intended to replace but must also have the required, possibly complex, 3-dimensional shape. As such, the replacement tissue is generally grown on or within a suitably shaped scaffold immersed in a culture medium in a bioreactor. The scaffold is a cell growth substrate which is shaped to provide growth of the tissue into the required 3-dimensional form. Within the bioreactor there is a (generally constant) flow of culture medium ensuring that the tissue-forming cells on or within the scaffold continuously receive a supply of nutrients and that the metabolic waste products of the cells are removed. The typically increased volumes of culture medium that can be used in bioreactors, compared to static culture flasks, allow immersion of scaffolds of a range of sizes suitable for production of a number of different tissue types. The perfusion of culture medium throughout the scaffold allows all cells to benefit from viable conditions for growth throughout the structure [1].

In addition, and particularly in the case of mechano-responsive tissue, it may be necessary to subject the tissue forming cells to mechanical stresses during their culture to produce fully functional tissue. Thus, for example, some types of

connective tissue such as bone, cartilage, ligament and tendons need to be subjected to mechanical stress during culturing thereof to give the required mechanical properties [2].

The degree of stress needed varies according to the cell types used and tissue types required. A number of ways of producing such stresses are known in the art, including direct mechanical stimulation of the cells, and hydrodynamic compression systems. The former method uses rollers or the like to compress the cells, whereas the latter utilises pulses of increased pressure in the culture medium supplying the bioreactor to mechanically stimulate the cells. None of the known methods of mechanically stimulating cells in order to produce functional tissue are, however, entirely satisfactory for many types of tissue such as bone, tendon and ligaments. Direct mechanical methods are cumbersome, and produce difficulties in maintaining the aseptic conditions required for culture. Hydrodynamic compression methods are generally ineffective. Moreover, all previously known methods suffer from the disadvantages that only one magnitude of stress can be applied across the range of cultured cells at any one time (generally a much higher stress than is required at a cellular level), and that the scaffold upon which the cells are grown must itself have considerable mechanical resilience to withstand the stresses applied to it.

It is therefore an object of the present invention to obviate or mitigate the above mentioned disadvantages.

According to a first aspect of the present invention there is provided a method of culturing tissue comprising growing tissue forming cells while subjecting the cells to mechanical stresses characterised in that the mechanical stresses are generated magnetically.

Thus in accordance with the invention, magnetically generated stresses are applied to the tissue forming cells so as to ensure that fully functional tissue is produced.

The method of the invention may be applied *in vitro* for the growth of tissue to be implanted in a patient. If effected *in vitro* it is preferred that the tissue forming cells are cultured on or in a 3-dimensional scaffold and preferably also in a bioreactor through which flows a tissue culture medium. Other types of tissue culture vessels may however be used. It is also possible for the method of the invention to be applied *in vivo* to grow new tissue *in situ* within the body of the a patient.

The stresses may be generated by a magnetic material capable of generating a force in response to a magnetic field applied within the bioreactor and transmitting that force to the tissue forming cells being cultured so as to apply the required stresses thereto. In preferred embodiments of the invention, the magnetic material is attached to the tissue forming cells and, for preference, takes the form of micro- or nano-particles, preferably coated magnetic micro- or nano-particles. Alternatively the magnetic material may be a ferrofluid which is inserted into the culture medium. A further possibility is the combined use of magnetic material attached to the cells and a ferrofluid.

Irrespective of the particular magnetic material used, the use of a time-varying magnetic gradient or homogeneous field modulates the movement of the magnetic material and consequently allows stresses to be repeatedly applied to the tissue forming cells. Such stresses may be accurately varied both in their magnitude and direction of application so that the tissue forming cells may be subjected to the required stress forming regime to ensure that fully functional tissue is produced. This can be accomplished by varying the magnetic properties of particles attached to different cells in different regions of the same scaffold (or different scaffolds) or through the use of the spatial variation of field strength in the gradient field.

The magnetic field may be varied at a frequency of, for example, 0.1 to 10 Hz. But, frequencies outside this range can also be used. The magnetic field will typically have a flux density on the order of (but not limited to) 10 mT to 1400 mT.

The magnitude of the stresses applied to the cells will be generally on the order of (but not limited to) 0.1 to 100 piconewtons (pN) and the direction in which the stress is applied may result from a linear, transitional motion (due to the gradient, particle does not need to be magnetically blocked) or rotational motion (due to the angle of the particle's magnetization vector with the applied field, must be a magnetically blocked particle) of the magnetic material in the applied magnetic field.

Significant advantages of the present invention are that (as indicated) it is easy to control the direction and magnitude of the applied stresses whilst maintaining aseptic conditions *in vitro* e.g. in a bioreactor or *in vivo* since variation of the magnetic field may be controlled remotely. Moreover the stresses that are generated at the cellular level are generally small (e.g. a few piconewtons) [3] and as such any scaffolds (on or in which the tissue forming cells are grown) do not need enhanced mechanical properties.

The method of the invention can be used to produce a variety of tissue types in both bioreactors and *in vivo* which require mechanical loading or activation of mechanosensitive ion channels. These include (but are not limited to) connective tissue such as bone, cartilage, ligament and tendons. Biopsies of the cells to be cultured may be obtained by standardized procedure [4].

It is also possible for the method of the invention to be applied to tissue constructs comprised of tissues of at least two different types, e.g. bone and cartilage. It is also possible to use human mesenchymal stem cells as a source which are differentiated into chondrocytes or bone cells *in situ* on or within scaffolds.

As mentioned above, a preferred embodiment of the invention involves attachment of magnetic micro- or nano- particles to the tissue forming cells for the purposes of applying the required stresses thereto. The magnetic micro- and nano-particles may be functionalised and attached to the tissue forming cells prior to seeding of the latter onto a scaffold on or in which the tissue is to be grown. Thus, for

example, the micro- and nano- particles may be coated with adhesion molecules, e.g. fibronectin and RGD molecules for attachment to the cells.

The micro- and nano- particles (intended to be attached to the cells) will generally be spherical or elliptical and have a diameter in the range 10nm to 10 μ m.

The particles for attachment to the cells may be coated or uncoated and single or multi-domain. Examples of suitable particles include, but are not limited to:

- (i) Coated magnetic microspheres ($d = 4 \mu\text{m}$) available from Spherotech, Inc. These microspheres consist of a magnetically blocked core –coated by a polymer.
- (ii) Single-domain, ferrite-doped silica nanoparticles with tunable size ($d = 50\text{--}300 \text{ nm}$) and narrow size distribution [5].

It is not however essential for the magnetic material to be particulate nor that it be attached to the cells. It is possible, for example, for the perfusion medium in a bioreactor or *in vivo* to contain a ferrofluid which is used to generate forces, from an applied magnetic gradient field, to the cells being cultured. The ferrofluid may for example be a PVA/magnetite nanoparticle-based ferrofluid ($d = 4\text{--}10\text{nm}$) [6]. It is possible also to use particles attached to the cells in combination with a ferrofluid.

The bioreactor may for example be a modification of an existing bioreactor such as perfusion, spinner flask, hydrodynamic compression and rotating vessel systems.

Conveniently the magnetic field is generated outside the tissue culture vessel (if the method is applied *in vitro*), or outside the body for the case of *in vivo* applications, and may be provided by a permanent magnet or an electromagnet. In order to generate variable fields, a permanent magnet may be moved relative to the cells being cultured. Thus, in the case of a bioreactor, such movement may for

example be longitudinally along the reactor, towards-and-away from the reactor or around the reactor. Any combination of these movements may also be used. In the case of an electromagnet, varying magnetic fields may be generated by provision of appropriate electric current levels to the electromagnet optionally in combination with movement of the electromagnet in the same manner as described for the permanent magnet.

Examples of commercially available magnets that may be used include Neodymium-Iron-Boron and Samarium-Cobalt permanent magnets that are capable of generating the required field gradients and flux densities. They can be geometrically tailored and magnetized to a variety of required specifications and produce flux densities at the surface in excess of 1 T (10,000 Gauss). Examples of electromagnets that may be used include cryo-cooled, superconducting magnetic coils capable of producing fields of several tesla.

The forces applied to the cells will generally be from 0.1 to 10pN (as previously indicated), such forces being capable of opening transmembrane ion channels. The magnetic fields and field gradients required to generate these forces vary depending on the magnetic, volumetric and shape properties of the particles and the distance between the tissue construct and the magnet. These parameters are governed by the equation:

$$F_{mag} = (\chi_2 - \chi_1) V \frac{1}{\mu_0} B(\nabla B)$$

where χ_2 is the volume magnetic susceptibility of the magnetic particle, χ_1 is the volume magnetic susceptibility of the surrounding medium (i.e. tissue/bone), μ_0 is the magnetic permeability of free space, B is the magnetic flux density in Tesla (T). Though this assumes spherical particles and no magnetic dipole interactions, it should give a good approximation of the field and gradient required for the system.

As the value of χ_1 for human tissue is very small and negative in comparison with the magnetic susceptibility of magnetite (or other magnetic material which will be used in the ferrofluids, and nanoparticles), χ_1 is negligible for this calculation and the expression $(\chi_2 - \chi_1)$ can be reduced to χ_2 . Also, as we are interested in the translational motion of the magnetite particle/fluid/material in an applied field along the z-axis (vertical) and, assuming a relative permeability of 1, the force expression can be reduced to:

$$F_{mag} = (\chi_2)VB \frac{dB}{dz}$$

for particles close to the magnetic field source.

It can be seen from these equations that the compressional (translational) force experienced by the tissue constructs in the presence of ferrofluids and magnetic particles is dependent on the strength of the field, the field gradient and the volumetric and magnetic properties of the particles. One of these parameters will have a strong spatial variation – the field strength/gradient product. This will enable the application of differential forces in three dimensions. In addition, by seeding different regions of the scaffold with particles, ferrofluids and magnetic materials of differing magnetic and volumetric properties, the three-dimensional variation in applied force can be enhanced. This facilitates the growth of complex tissue structures via the spatial variation of applied forces inside the bioreactor.

A number of different scaffold types (essentially 3-dimensional porous blocks which can be varied in dimension) can be used. One of the advantages of the present invention is that the scaffolds do not need enhanced mechanical properties as the stresses involved are generally small. It is possible, for example, to use a biodegradable, porous polylactic acid (PLA) based scaffold. Alternative scaffolds include PGA (poly glycolic acid) materials which are rapidly degrading and are less mechanically strong and collagen scaffolds which are natural materials [7]. The

scaffold may be coated with collagen type 1 or other adhesion molecules (such as RGD or non-RGD based molecules) to improve cell adhesion.

The invention will be illustrated, by way of example only, with reference to the accompanying drawings, in which:

Fig 1 illustrates a first embodiment of the invention;

Fig 2 illustrates a second embodiment of the invention; and

Fig 3 illustrates activation of a mechano-sensitive transmembrane ion channel by the use of a magnetic field.

Referring to Fig 1, there is illustrated a tubular bioreactor 1 associated with permanent magnets 2 which are positioned externally of the reactor 1 and which are mounted on a carrier arrangement 3 connected to a computer-controlled (or other time-varying) drive system not illustrated in detail in the drawings.

Within the bioreactor 1 are a number of longitudinally spaced tissue constructs 4 each depicted as being comprised on bone tissue on one side (the side remote from the magnets 3) and cartilage tissue on the other side (other tissue types may also be used). The tissue cells have magnetic beads (not shown) attached thereto and are seeded on 3-D scaffolds (again not shown). Nutrients are supplied to the bioreactor as depicted by arrow 5 and exit therefrom as depicted by arrow 6.

There are a total of four magnets 2 (though this number can vary to match the number of tissue constructs) which are external of the bioreactor and longitudinally spaced therealong. The positioning of the magnets is such that there is one magnet associated with each of the tissue constructs, the magnets being provided on the cartilage sides thereof.

In use of the apparatus, the carrier is driven so as to oscillate the magnets transversely towards and away from the bioreactor. The oscillation frequency at which the magnets are driven will usually be varied and generally be in the range of 0.1 to 10 Hz although values outside this range may be used.

The oscillation of the magnets stimulates a compression/relaxation cycle which is applied to the tissue constructs, the frequency of which can also be varied by mechanical drivers (not shown) attached to the magnets. The magnet field gradient (spatially varying magnetic field strength) ensures that the cartilage experiences slightly higher flux densities than the bone cells.

Strong magnetic field gradients will produce a translational motion on the nanoparticles directed towards the magnets, compressing the cells and scaffold inside the bioreactor. This compression will simulate mechanical loading without requiring direct access to the cells inside the bioreactor. Loads can be easily varied by changing the magnetic field strength and gradient, magnet position and/or the physical properties of the nanoparticles compressing the tissue constructs.

If desired, the magnetic particles associated with the bone cells may have different magnetic properties from those associated with the cartilage so that different mechanically stresses are applied to the two different types of cells.

Fig 2 illustrates a modification of the apparatus shown in Fig 1. In the modification of Fig 2, the (permanent) magnets are oscillated parallel to the longitudinal axis of the bioreactor (rather than transversely to the axis in the case of Fig 1).

A number of modifications may be made to the illustrated embodiments.

Thus, for example, the magnets may be swept relatively around the bioreactor. This may most conveniently, but not necessarily, be achieved by keeping the magnets fixed and rotating the bioreactor around its longitudinal axis.

Alternatively or additionally the permanent magnets illustrated in Figs 1 and 2 may be replaced by electromagnets. A further possibility is for the nanoparticles attached to the cells to be replaced by ferrofluids. If desired, a combination of attached nanoparticles and ferrofluids may also be used.

A further possibility is to use magnetic/metal plates or other structures which could be attracted to the magnets in order to deform the entire scaffold.

Reference is now made to Fig 3 which illustrates an alternative method of activation of mechano-sensitive transmembrane ion channels by the use of a magnetic field so as to simulate period mechanical loading of a tissue construct.

More particularly, Fig 3 illustrates a cell 10 having a membrane 11 enclosing the cell's cytoplasm 12. Within membrane 11 is a mechanosensitive ion channel 13. A functionalized magnetically blocked particle 14 (such as Sphereotech's coated ferromagnetic particles, $d=4.5\mu\text{m}$) is rigidly attached to the cell membrane 11 either directly or indirectly via cytoskeletal coupling [8, 9].

In the condition shown in Fig 3(a), no magnetic field is applied to the cell 10 and the ion channel 13 is closed.

By oscillating a magnetic field source (not shown) the magnetic particle 14 attached to the cell 10 can be twisted, exerting a mechanical stress on the cell membrane 11 and activating the mechano-sensitive ion channel 13 (Figure 3b). This ion channel activation initiates biochemical reaction pathways in the cells being cultured and simulates periodic mechanical loading of tissue constructs inside the bioreactor.

REFERENCES:

1. Ying, Y., Peak, M., Magnay, J., and El Haj, A.J. (2000) Dynamic cell scaffold interactions: implications for tissue engineering. *Proceedings of the second Smith and Nephew international symposium on tissue engineering, York, UK.*
2. El Haj, AJ, LM Walker, MR Preston, SJ Publicover (1999) Mechanotransduction pathways in bone: calcium fluxes and the role of voltage-operated calcium channels. *Med. Biol. Eng. Comp.*, 37: 403-409.
3. Howard J and AJ Hudspeth (1989) Compliance of the hair bundle associated with the gating of mechanoelectrical transduction channels in the bullfrog's saccular hair cell. *Neuron* 1: 189-199.
4. Walker et al J Cell Biochem 2000.
5. Pardoe, H, W Chua-anusorn, TG St. Pierre, J Dobson (2000) Structural and magnetic properties of nanoscale magnetic particles synthesised by coprecipitation of iron oxide in the presence of dextran or polyvinyl alcohol. *J. Magn. Mag. Materials*. In Press.
6. Tan, W, S Santra, Z Peng, R Tapecc and J Dobson (2000) Coated nanoparticles. *US Patent Pending* (Filed May 17, 2000).
7. Sittinger et al 1996.
8. Kirschvink, JL (1992) Comments on "Constraints on biological effects of weak extremely-low-frequency electromagnetic fields". *Phys. Rev. A*. 46: 2178-2184.
9. Dobson, J and TG St. Pierre (1996) Application of the Ferromagnetic Transduction Model to D.C. and Pulsed Magnetic Fields: Effects on Epileptogenic Tissue and Implications for Cellular Phone Safety. *Biochem. Biophys. Res. Commun.*, 227:718-723.

CLAIMS

1. A method of culturing tissue comprising growing tissue forming cells whilst subjecting tissue forming cells to mechanical stresses characterised in that the mechanical stresses are generated magnetically.
2. A method as claimed in claim 1 wherein the stresses are generated by a magnetic material capable of generating a force in response to a magnetic field applied and transmitting that force to the tissue forming cells.
3. A method as claimed in claim 2 wherein the magnetic material is attached to the tissue forming cells.
4. A method as claimed in claim 3 wherein the magnetic material comprises micro- or nano-particles.
5. A method as claimed in any one of claims 2 to 4 wherein the magnetic material comprises a ferrofluid.
6. A method as claimed in any of claims 1 to 5 wherein the magnetic field is varied during culture of the these tissue forming cells.
7. A method as claimed in claim 6 wherein the magnetic field is varied sinusoidally.
8. A method as claimed in claim 6 or 7 wherein the magnetic field is varied at a frequency of 0.1 to 10Hz.
9. A method as claimed in any one of claims 1 to 8 which is applied *in vitro*.
10. A method as claimed in claim 9 wherein the tissue forming cells are grown on or in a 3-dimensional scaffold.

11. A method as claimed in claim 9 or 10 wherein the tissue forming cells are cultured in a bioreactor through which flows a culture medium.
12. A method as claimed in claim 11 wherein the magnetic field is applied from externally of the bioreactor.
13. A method as claimed in any one of claims 1 to 8 applied *in vivo*.
14. A method as claimed in any one of claims 1 to 13 wherein the cells being cultured are for forming connective tissue.
15. A method as claimed in claim 14 wherein the cells being cultured are for forming bone, cartilage, ligament or tendons.
16. A method as claimed in any one of claims 1 to 15 wherein two or more different cell types are cultured.
17. A method as claimed in claim 16 wherein different stresses are applied to the different cell types.
18. A method as claimed in claim 16 or 17 wherein the cells being cultured comprise bone-forming cells and cartilage-forming cells.
19. A method as claimed in any one of claims 1 to 18 wherein the forces applied to the cells or scaffold are in the range 0.01 to 10 Newtons.
20. A bioreactor having an inlet and an outlet for a culture medium and being provided means for applying a variable magnetic field within the bioreactor.
21. A bioreactor as claimed in claim 20 wherein the means for applying the variable magnetic field comprise an electromagnet.

22. A bioreactor as claimed in claim 21 wherein the means for applying the variable magnetic field comprises at least one permanent magnet which is moveable relative to the bioreactor.




FIG. 1

2 / 3





FIG. 2



(19) World Intellectual Property Organization
International Bureau




(43) International Publication Date
4 July 2002 (04.07.2002)

PCT

(10) International Publication Number
WO 02/051985 A3

- (51) International Patent Classification⁷: C12N 5/08, C12M 3/00 (74) Agent: HARRISON GODDARD FOOTE, 31 St. Saviourgate, York YO1 8NQ (GB).
- (21) International Application Number: PCT/GB01/05606 (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.
- (22) International Filing Date:
19 December 2001 (19.12.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
0031462.5 22 December 2000 (22.12.2000) GB
0031651.3 23 December 2000 (23.12.2000) GB
60/267,032 7 February 2001 (07.02.2001) US
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (for all designated States except US): KEELE UNIVERSITY [GB/GB]; Keele, Staffordshire ST5 5BG (GB).
- Published:
— with international search report
- (72) Inventors; and
(75) Inventors/Applicants (for US only): EL-HAJ, Alicia, Jennifer, Hafeeza [GB]; Beardhough Cottage, White Road, New Mills, High Peak SK22 4EQ (GB). DOBSON, Jon, Paul [GB]; Claremount Brook Lane, Endon, stoke-on-Trent ST9 9EZ (GB).
- (88) Date of publication of the international search report:
16 January 2003
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: CULTURING TISSUE USING MAGNETICALLY GENERATED MECHANICAL STRESSES



(57) Abstract: A method of culturing tissue comprises growing tissue forming cells whilst subjecting the tissue forming cells to mechanical stresses which are generated magnetically.

WO 02/051985 A3

INTERNATIONAL SEARCH REPORT

Application No

GB 01/05606

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C12N5/08 C12M3/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C12M

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	YUGE LOUIS ET AL: "Differentiation of myoblasts is accelerated in culture in a magnetic field." IN VITRO CELLULAR & DEVELOPMENTAL BIOLOGY ANIMAL, vol. 36, no. 6, June 2000 (2000-06), pages 383-386, XP001061490 ISSN: 1071-2690 page 383, right-hand column; figure 1 page 385 --- -/--	1,2,4,9, 12



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

21 June 2002

Date of mailing of the international search report

06.09.02

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Steffen, P

INTERNATIONAL SEARCH REPORT

Int. Application No.
PCT/GB 01/05606

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	LIU MINGYAO ET AL: "Bio-stretch, a computerized cell strain apparatus for three-dimensional organotypic cultures." IN VITRO CELLULAR & DEVELOPMENTAL BIOLOGY ANIMAL, vol. 35, no. 2, February 1999 (1999-02), pages 87-93, XP001080069 ISSN: 1071-2690 page 87, right-hand column, last paragraph -page 88, left-hand column, paragraph 1; figures 1,3 page 89 -page 92	1,2,6, 8-10,12
X	KASPAR D ET AL: "Dynamic cell stretching increases human osteoblast proliferation and CICP synthesis but decreases osteocalcin synthesis and alkaline phosphatase activity." JOURNAL OF BIOMECHANICS, vol. 33, no. 1, January 2000 (2000-01), pages 45-51, XP002203098 ISSN: 0021-9290 page 46; table 1 page 48; figure 2	1,2,6-9, 12,14, 15,19
A	-& BOTTLANG M ET AL: "A cell strain system for small homogeneous strain applications." BIOMEDIZINISCHE TECHNIK, vol. 42, no. 11, November 1997 (1997-11), pages 305-309, XP001080015 ISSN: 0013-5585 page 306	

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

Application No

GB 01/05606

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>GLOGAUER MICHAEL ET AL: "A new method for application of force to cells via ferric oxide beads."</p> <p>PFLUEGERS ARCHIV EUROPEAN JOURNAL OF PHYSIOLOGY, vol. 435, no. 2, January 1998 (1998-01), pages 320-327, XP002203099 ISSN: 0031-6768 page 320, right-hand column -page 321, right-hand column, paragraph 2; figure 1 page 323 page 324, left-hand column, paragraph 2 -page 325, left-hand column, paragraph 2 -& DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US1995 GLOGAUER, M. ET AL.: "Magnetic fields applied to collagen-coated ferric oxide beads induce stretch-activated Ca-2+ flux in fibroblasts." Database accession no. PREV199698605117 XP002203103 abstract & AMERICAN JOURNAL OF PHYSIOLOGY, vol. 269, no. 5 PART 1, 1995, pages C1093-C1104, ISSN: 0002-9513</p>	1-4, 9, 12, 19
X	<p>SMALT R ET AL: "Induction of NO and prostaglandin E2 in osteoblasts by wall-shear stress but not mechanical strain." AMERICAN JOURNAL OF PHYSIOLOGY, vol. 273, no. 4 PART 1, October 1997 (1997-10), pages E751-E758, XP002203100 ISSN: 0002-9513 page E752; figure 1</p>	1, 2, 6-8, 12, 14, 15
X	<p>US 5 486 457 A (BUTLER JAMES P ET AL) 23 January 1996 (1996-01-23) figure 1; examples 1, 2</p>	1-4
X	<p>WANG NING ET AL: "Contribution of intermediate filaments to cell stiffness, stiffening, and growth." AMERICAN JOURNAL OF PHYSIOLOGY, vol. 279, no. 1 Part 1, July 2000 (2000-07), pages C188-C194, XP002203101 ISSN: 0002-9513 page C189, left-hand column</p> <p style="text-align: center;">-/--</p>	1-4

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

Int. Application No.

PC1/GB 01/05606

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WANG NING: "Mechanical interactions among cytoskeletal filaments." HYPERTENSION (DALLAS), vol. 32, no. 1, July 1998 (1998-07), pages 162-165, XP002203102 ISSN: 0194-911X page 162, right-hand column, paragraph 2 -page 163, left-hand column, paragraph 1 -----	1-4

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

International application No.
PCT/GB 01/05606

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.

2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.

3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1-19

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-19

A method of culturing tissue comprising growing tissue forming cells whilst subjecting tissue forming cells to mechanical stresses characterised in that the mechanical stresses are generated magnetically. Furthermore further embodiments of said method.

2. Claims: 20-22

A bioreactor having an inlet and an outlet for a culture medium and being provided with means for applying a variable magnetic field within the bioreactor. Furthermore further embodiments of said bioreactor.

INTERNATIONAL SEARCH REPORT

Information on patent family members

1st Application No

PCT/GB 01/05606

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5486457 A	23-01-1996	WO 9506248 A	02-03-1995

Form PCT/ISA/210 (patent family annex) (July 1992)